

TITLE OF THE INVENTION

GREEN FLUORESCENT PROTEINS AND BLUE FLUORESCENT
PROTEINS

BACKGROUND OF THE INVENTION5 Field of the invention

This invention relates to novel fluorescent
proteins, GFPs and BFPs.

Related background art

10 GFP (Green Fluorescent Protein), which was found
in *Aequorea victoria*, is a relatively small protein
having a molecular weight of 26,900 and comprising
the overall 238 amino acid residues as shown below
(SEQ No. 1 in the Sequence Listing).

15 Met Ser Lys Gly Glu Glu Leu Phe Thr Gly Val Val Pro Ile Leu Val
1 5 10 15
Glu Leu Asp Gly Asp Val Asn Gly His Lys Phe Ser Val Ser Gly Glu
20 25 30
Gly Glu Gly Asp Ala Thr Tyr Gly Lys Leu Thr Leu Lys Phe Ile Cys
20 35 40 45
Thr Thr Gly Lys Leu Pro Val Pro Trp Pro Thr Leu Val Thr Thr Phe
50 55 60
Ser Tyr Gly Val Gln Cys Phe Ser Arg Tyr Pro Asp His Met Lys Gln
70 75 80
25 His Asp Phe Phe Lys Ser Ala Met Pro Glu Gly Tyr Val Gln Glu Arg
85 90 95

Thr Ile Phe Phe Lys Asp Asp Gly Asn Tyr Lys Thr Arg Ala Glu Val

100

105

110

Lys Phe Glu Gly Asp Thr Leu Val Asn Arg Ile Glu Leu Lys Gly Ile

115

120

125

5 Asp Phe Lys Glu Asp Gly Asn Ile Leu Gly His Lys Leu Glu Tyr Asn

130

135

140

Tyr Asn Ser His Asn Val Tyr Ile Met Ala Asp Lys Gln Lys Asn Gly

145

150

155

160

Ile Lys Val Asn Phe Lys Ile Arg His Asn Ile Glu Asp Gly Ser Val

10

165

170

175

Gln Leu Ala Asp His Tyr Gln Gln Asn Thr Pro Ile Gly Asp Gly Pro

180

185

190

Val Leu Leu Pro Asp Asn His Tyr Leu Ser Thr Gln Ser Ala Leu Ser

195

200

205

15 Lys Asp Pro Asn Glu Lys Arg Asp His Met Val Leu Leu Glu Phe Val

210

215

230

Thr Ala Ala Gly Ile Thr His Gly Met Asp Glu Leu Tyr Lys

225

230

235

238

20 In the present specification, the term "GFP
protein" refers to a protein that emits green
fluorescence when excited by ultraviolet-blue light
and that, then, does not require an energy source
such as a special substrate or ATP. In other words,
25 the chromophore formation reaction of GFP is
autonomous, and the portion of serine-tyrosine-

glycine at Nos. 65-67 from the amino terminus forms an imidazolidine ring oxidatively which serves as a chromophore. (Yuichiro Watanabe, Gendai Kagaku "Modern Chemistry" 12, 46-52 (1995); R. Heim et al. Proc. Natl. Acad. Sci. USA 91: 12501-12504 (1994).) Because GFP possesses such a property, a DNA encoding this protein is linked to a suitable expression vector and is introduced into the desired cells to express GFP, which alone results in fluorescent images. Therefore, GFP is in use for the visual analysis of gene expression and localization of proteins in a variety of cells in their viable state. However, since such GFP was not luminous at 37 °C, there was a problem that culturing must necessarily be done at 30 °C for the purpose of observation in mammalian cells or the like. In connection with this problem, it has been reported that the mutations of V163A and S175G enhance the thermal stability. (K. R. Siemering et al. Curr. Biol. 6, 1653-1663 (1996).)

Recently, a mutant of GFP into which the mutations of Y66H and Y145F were introduced and which had different wavelength characteristics (it is also referred to as "Mutant," and its amino acid sequence is described below with the above-mentioned mutations shown as underlined) was developed. This is referred to as "BFP (Blue Fluorescent Protein)," because it

emits blue fluorescence by UV excitation. (R. Heim et al. Curr. Biol. 6, 178-182 (1996); R. Heim et al. Proc. Natl. Acad. Sci. USA 91, 12501-12504 (1994).)

In the present specification, the term "BFP protein" refers to a protein that emits blue fluorescence when excited by ultraviolet-blue light and that, then, does not require an energy source such as a special substrate or ATP. However, such BFP had a problem that it experienced severe fading as compared to GFP and was difficult to be observed under a microscope or the like. As used herein to designate mutation, the position of the mutation is expressed by a specific amino acid number in the sequence of the above-mentioned wild type; the amino acid prior to its mutation is described preceding the number and the mutated amino acid is to be described following the number.

Further, amino acids are designated by the one-letter code or three-letter code as appropriate.

Met Ser Lys Gly Glu Glu Leu Phe Thr Gly Val Val Pro Ile Leu Val

1

5

10

15

Glu Leu Asp Gly Asp Val Asn Gly His Lys Phe Ser Val Ser Gly Glu

20

25

30

Gly Glu Gly Asp Ala Thr Tyr Gly Lys Leu Thr Leu Lys Phe Ile Cys

35

40

45

B

5

The present invention will become more fully understood from the detailed description given hereinbelow and the accompanying drawings which are 5 given by way of illustration only, and thus are not to be considered as limiting the present invention.

1a2 In view of the above-mentioned problems, the present inventors made extensive researches and succeeded in the discovery of novel GFPs and BFPs that are free from such problems by introducing certain mutations into specific positions of the amino acid sequence for GFP or BFP, thus accomplishing this invention.

6

5

15

20

- 25

Furthermore, on the basis of the amino acid sequence for the wild type of BFP as described above, GFPs into which the mutations as described below had been introduced were prepared, and their fluorescence and thermal characteristics were investigated in this invention. Here, the mutations introduced were based on the amino acid sequence for the wild type of GFP.

(4) Y66H, Y145F: Phe64Leu, Leu236Arg

(5) Y66H, Y145F: Phe64Leu

(6) Y66H, Y145F: Val163Ala, Ser175Gly

(7) Y66H, Y145F: Phe64Leu, Val163Ala, Ser175Gly, Leu236Arg

Consequently, it was discovered that the resulting BFP and GFP mutants had improved fluorescence characteristics and thermal stability. Specifically, this invention provides novel BFPs and GFPs as will be described below, and further, genes coding them.

1. A GFP protein comprising the amino acid sequence set forth in SEQ ID No. 1 in the Sequence Listing, said sequence having at least mutations of Phe64Leu, Val163Ala, and Ser175Gly.

2. A GFP protein comprising the amino acid sequence set forth in SEQ ID No. 1 in the Sequence Listing, said sequence having the three mutations of Phe64Leu, Val163Ala, and Ser175Gly.

3. A BFP protein comprising the amino acid

sequence set forth in SEQ ID No. 1 in the Sequence Listing, said sequence having at least mutations of Y66H, Y145F, and Phe64Leu.

5 4. A BFP protein comprising the amino acid sequence set forth in SEQ ID No. 1 in the Sequence Listing, said sequence having at least mutations of Y66H, Y145F, Phe64Leu, and Leu236Arg.

10 5. A BFP protein comprising the amino acid sequence set forth in SEQ ID No. 1 in the Sequence Listing, said sequence having the four mutations of Y66H, Y145F, Phe64Leu, and Leu236Arg.

15 6. A BFP protein comprising the amino acid sequence set forth in SEQ ID No. 1 in the Sequence Listing, said sequence having at least mutations of Y66H, Y145F, Phe64Leu, Val163Ala, Ser175Gly and Leu236Arg.

20 7. A BFP protein comprising the amino acid sequence set forth in SEQ ID No. 1 in the Sequence Listing, said sequence having the six mutations of Y66H, Y145F, Phe64Leu, Val163Ala, Ser175Gly and Leu236Arg.

8. A gene encoding the GFP protein according to either Item 1 or Item 2 as described above.

25 9. A gene encoding the BFP protein according to any of Items 3-7 as described above

This invention will be illustrated in detail

hereinbelow based on its embodiments. The abbreviations of nucleic acids and amino acids (one-letter and three-letter codes) as used in the present specification are set forth below.

5	(Nucleic Acids)	
	DNA	deoxyribonucleic acid
	A	Adenine
	C	Cytosine
	G	Guanine
10	T	Thymine
	(Amino Acids)	
	Ala (A)	Alanine
	Arg (R)	Arginine
	Asn (N)	Asparagine
15	Asp (D)	Aspartic acid
	Cys (C)	Cysteine
	Gln (Q)	Glutamine
	Glu (E)	Glutamic acid
	Gly (G)	Glycine
20	His (H)	Histidine
	Ile (I)	Isoleucine
	Leu (L)	Leucine
	Lys (K)	Lysine
	Met (M)	Methionine
25	Phe (F)	Phenylalanine
	Pro (P)	Proline

Ser (S)	Serine
Thr (T)	Threonine
Trp (W)	Tryptophan
Tyr (Y)	Tyrosine
Val (V)	Valine

5

BRIEF DESCRIPTION OF THE DRAWINGS

10

The file of this patent contains at least one drawing executed in color. Copies of this patent with color drawing(s) will be provided by the Patent and Trademark Office upon request and payment of the necessary fee.

15

FIG. 1 is an electrophoresis photograph obtained when *E. coli* harboring each plasmid was induced by IPTG and its protein was subjected to SDS-PAGE, where Lanes 1, 4, and 7 show the results of electrophoresis of 50 μ l equivalents of the *E. coli* culture media and Lanes 2, 3, 5, 6, 8, and 9 show those of 50 μ l equivalents.

20

FIG. 2 is a photograph showing the fluorescence emitted when the *E. coli* harboring each plasmid was streaked on a plate, and after culturing at 37 °C overnight, it was irradiated with UV at a long wavelength. "1" indicates empty vector(pQE30), "2" indicates original GFP(pQE101). "3" indicates improved GFP(F64L/V163A/S175G)(pQE105). "4" indicates original BFP(pQE201). "5" indicates improved

25

BFP(F64L/L236R)(pQE202). "6" indicates improved
BFP(F64L/V163A/S175G/L236R)(pQE205).

FIG. 3 is an electrophoresis photograph showing
the results obtained when CHO cells, after
transfection with each plasmid, were cultured at
37 °C or at 30 °C, and the culture was subjected to
SDS-PAGE followed by transfer onto a nitrocellulose
membrane and western blotting with an anti-GFP
antibody. Here, the arrow indicates GFP or BFP.

FIGS. 4A and 4B are fluorescence photographs
showing the effects associated with the improved type
of GFP(B) with the original type of GFP(A).

FIGS. 4C and 4D are fluorescence photographs
showing the effects associated with the improved type
of BFP(D) with the original type of BFP(C).

~~DESCRIPTION OF THE PREFERRED EMBODIMENTS~~

Novel GFP or BFP proteins according to this
invention are those obtained by introducing certain
mutations to parts of the amino acid sequences for
the wild types of GFP and BFP, and exhibit improved
fluorescence characteristics and thermal stability.
Therefore, this invention embraces proteins having at
least such amino acid sequences insofar as they
exhibit the improved fluorescence characteristics and
thermal stability based on the novel GFP or BFP
proteins according to the invention. Namely, in the

00852000-051001

INSA3

123

cases where cells of a variety of origins are used as will be in use in the Examples below, the invention also embraces proteins to which a variety of amino acid sequences other than the aforementioned amino acid sequences are appended at their N- or C-termini and which exhibit the improved fluorescence characteristics and thermal stability based on the novel GFP or BFP proteins according to the invention.

Moreover, this invention provides genes encoding such novel proteins or proteins containing them within parts thereof.

There are no particular limitations to methods for obtaining the novel GFPs or BFPs according to this invention, and methods for artificially obtaining them by means of chemical syntheses and methods for obtaining them according to standard genetic engineering are possible. The latter methods are made possible through the genetic engineering techniques in which suitable vectors conventionally known and means for introducing mutations are combined. Concretely, the following procedure is preferred.

Specifically, the procedure comprises the steps of: (1) starting with a known GFP or BFP protein to be improved and introducing a gene encoding said protein into a suitable vector; (2) introducing

mutations into said gene selectively or randomly according to known methods; and (3) selecting desirable mutants on the basis of the fluorescence intensities and temperature-dependence, among others, of the resultant GFP or BFP mutants.

The Contents of Application No.026418/1998, filed on January 23, 1998 in Japan is hereby incorporated by reference.

The above-mentioned procedure will be hereinbelow illustrated in detail by way of examples; however, this invention is not to be limited to these specific examples.

EXAMPLES

(I) The genetic engineering techniques as used in the present examples will be illustrated in the following.

1. Vector Construction

In this invention, a DNA portion encoding GFP of pGFP-C1 vector (available from Clontech Inc.) was replaced by a DNA of GFP derived from phGFP-S65T (available from Clontech Inc.), which served as a basic plasmid (hereinafter referred to as "phGFP(101)-C1"). The vector is meant for expression in mammalian cells and its full base sequence including the vector part is known in the art. The corresponding amino acid sequence is set forth below.

15

Lys Asp Pro Asn Glu Lys Arg Asp His Met Val Leu Leu Glu Phe Val

210

215

230

Thr Ala Ala Gly Ile Thr Leu Gly Met Asp Glu Leu Tyr Lys

225

230

235

238

5

10

15

20

25

Here, the protein encoded by phGFP-S65T as described above is compared with that of a wild type derived from jellyfish: (i) an amino acid (valine) has been inserted between methionine, which is amino acid number 1 of the amino acid sequence, and serine, which is amino acid number 2; (ii) serine, which is amino acid number 65, has been further mutated to threonine; and (iii) histidine, which is amino acid number 231, has been mutated to leucine. These are respectively underlined in the amino acid sequence as described above. Thus for example, the amino acid number 65 threonine becomes number 66 in reality, but amino acid sequence numbers corresponding to those of the wild type are employed for the amino acid numbers connected with mutation, in accordance with general rules. In other words, the amino acid numbers for the amino acid sequence of the wild type derived from jellyfish (amino acid numbers 1 through 238) are to be used. The extra valine as described above is construed as having been inserted between amino acid number 1 and amino acid number 2, and no number is

then designated therefor. In practice, such an addition of valine has been used as a working example to illustrate the embodiments of this invention and it is not the essential amino acid sequence of this invention. Accordingly, in the explanation that follows the presence (or the absence) of the valine addition will not affect the scope of the invention.

Furthermore, methods for introducing specific mutations are not particularly limited, and for example, the method of introduction used in the examples of this invention as described below is feasible. Specifically, a DNA region encoding GFP was cut out from the above-mentioned pHGFP(101)-C1 with HindIII, and it was inserted into the HindIII site of a pUC18 vector or a pQE30 vector (Qiagen) to thereby prepare pUCGFP(101) or pQEGFP(101). Here, the pQE30 vector was meant for expression in *E. coli*.

Employing the resultant pUCGFP(101), pUCBFP(101) into which the mutations of T65S, Y66H, and Y145F had been introduced by the site-directed mutation introduction method as described below was prepared.

Here, through said mutation the amino acid number 65 Ser that was introduced by the above-mentioned mutation (T65S) proved to be identical with the wild type site.

Further, a DNA encoding BFP was cut out from the

obtained pUCBFP(201) by digestion with EcoRI/XhoI and it was cloned into the EcoRI/XhoI site of Bluescript II KS(-) (Stratagene) to thereby prepare blueBFP(201).

Furthermore, a DNA region encoding BFP was cut out from the obtained pUCBFP(201) by digestion with HindIII and it was inserted into the HindIII site of a pQE30 vector to thereby prepare pQEBFP(201). On the other hand, phBFP(201)-Cl was prepared by replacing the GFP coding region of the phGFP(101)-Cl vector with the above-mentioned DNA in like manner.

2. Mutagenic Polymerase Chain Reaction (hereinafter referred to as "PCR")

INS A4 ^{1a4} ~~Moreover, methods for randomly introducing mutations are not particularly limited, and Mutagenic PCR as described below can preferably be used in this invention. The Mutagenic PCR can be carried out according to methods known in the art. (C. W. Dieffenbach, ed. PCR PRIMER, A Laboratory Manual (Cold Spring Harbor Laboratory Press) (1955) pp. 583-588.)~~ Concretely, the following conditions were employed in the examples.

INS A5 ^{1a5} ~~About 50 ng of Plasmid BlueBFP(201) was added to 10xmutagenic PCR buffer (70 mM MgCl₂, 500 mM KCl, and 100 mM Tris-HCl, pH 8.3 at 25 °C; 0.1% (w/v) gelatin) 10 μl, 10x dNTP (2 mM dGTP, 2 mM dATP, 10 mM dCTP, and 10 mM dTTP) 10 μl, 10 pmol/μl primer (23mer~~

09852000-051001
M13Universal primer and M13Reverse primer) 3 μ l, and
a H₂O 62 μ l, and mixed. Subsequently, 10 μ l of 5 mM
MnCl₂ was added and mixed, and 1 μ l of Taq Polymerase
(Takara) was added to conduct PCR (PC-700 available
5 from ASTEC Inc. was used). The PCR was conducted in
three tubes under the following conditions: 25 cycles
a at 94 °C for 1 min, 30 cycles at 45 °C for 1 min, and
35 cycles at 72 °C for 1 min, respectively.

10 After the respective reaction solutions were
combined and treated with chloroform twice, a DNA
fragment encoding the amplified BFP was recovered by
carrying out electrophoresis on a 1% agarose gel
after digestion with BamHI and XhoI and it was
inserted into the BamHI and SalI sites of pQE30
15 (Qiagen Inc.).

Transformation was performed on *E. coli* JM109,
and inoculation was done in a LB medium containing
carbenicillin to incubate JM109 at 37 °C for 16 h.
Subsequently, the incubated product was allowed to
20 stand at room temperature for 24 h. The *E. coli*
colonies that resulted on a plate were irradiated
with UV (Funakoshi UV Transilluminator FTI-201 UV 365
nm) from the top side of the plate for 1 h, and
colonies emitting sufficient illumination visually
25 after irradiation were selected: ten colonies were
obtained in the example.

Sequence determination was performed on the selected plasmids. With respect to the mutant having mutations within its coding region that appeared meaningful, the coding region was cut out with Hind3 and was inserted into the HindIII site of pQE30, and thereafter, this was cut out with SalI/BglII and replaced by the corresponding portion of pQEBFP to bring the cloning site of the vector into conformity with pQEBFP(201): in the present examples the one prepared from Mutant 10 was designated pQEBFP (202).

3. Construction of Mutant GFP/BFP by the Site-Directed Mutation Introduction Method

The site-directed mutation introduction methods are not particularly limited, and for example, the protocol for a Quick Change Kit from Stratagene Inc. was followed. The oligonucleotides shown in Table 2 below were used as primers and the plasmid (about 0.03 μ g) obtained by subcloning GFP or BFPcDNA into the HindIII site of a pUC18 or pQE30 vector was used as a template. The concrete PCR conditions are preferably as follows: 16 cycles at 95 °C for 30 sec, 55 °C for 1 min, and 68 °C for 10 min.

TABLE 2

oligo no.		sequence	
5	1F	TCGTGACCACCTTCTCCCACGGCCTGCA	
	1R	TGCACGCCGTGGGAGAAGGTGGTCACGA	
	2F	GCTGGAGTACAACCTCAACAGCCACAACG	
	2R	CGTTGTGGCTGTTGAAGTTGTACTCCAGC	
	3F	CCTCGTGACCACCCTCTCCCACGGCGTG	
10	3R	CACGCCGTGGGAGAGGGTGGTCACGAGG	
	4F	CCTCGTGACCACCCTCACCTACGGCGTG	
	4R	CACGCCGTAGGTGAGGGTGGTCACGAGG	
	5F	GAACGGCATCAAGGCCAACTTCAAGATCC	
	5R	GGATCTTGAAGTTGGCCTTGATGCCGTTT	
15	6F	CATCGAGGACGGCGGCGTGCAGCTCGCC	
	6R	GGCGAGCTGCACGCCGCCGTCCTCGATG	

09852000-051001

TABLE 3

	GFP or BFP used as template	oligo no. used in the introduction of mutation	GFP mutant or BFP mutant after mutation-introduction
5	pUCGFP(101)	1F+1R	pUCGFP101(+Y66H)
	pUCGFP101(+Y66H)	2F+2R	pUC(201)
	pQEGFP(101)	4F+4R	pQEGFP(103)
10	pQEBFP(201)	3F+3R	pQEBFP(203)
	pQEGFP(101)	5F+5R	pQEGFP101(+V163A)
	pQEGFP101(+V163A)	6F+6R	pQEGFP(104)
	pQEBFP(201)	5F+5R	pQEBFP201(+V163A)
	pQEBFP201(+V163A)	6F+6R	pQEBFP(204)
15	pQEGFP(104)	4F+4R	pQEGFP(105)
	pQEBFP(202)	5F+5R	pQEBFP202(+V163A)
	pQEBFP202(+V163R)	6F+6R	pQEBFP(205)

20 Sequence determination of the resulting plasmids was conducted and it was verified that the desired mutations were contained in the plasmids.

25 In the examples of this invention, GFPs were designated as 101-105 and BFP were designated as 201-205 for reasons of convenience to place a variety of mutants as obtained in good order. Table 4 below

thus summarizes the mutations introduced. Although not shown in the table, GFP 101-105 all contain the mutations of Ser65Thr and His231Leu.

5 ~~TABLE 4~~

1a6

GFP

101 none

103 Phe64Leu

104 Val163Ala, Ser175Gly

105 Phe64Leu, Val163Ala, Ser175Gly

BFP (as for BFP, the two mutations, Y66H and Y145F, have been introduced into the sequence for GFP which serves as a basis)

201 Y66H, Y145F

202 Y66H, Y145F: Phe64Leu, Leu236Arg

203 Y66H, Y145F: Phe64Leu

204 Y66H, Y145F: Val163Ala, Ser175Gly

205 Y66H, Y145F: Phe64Leu, Val163Ala, Ser175Gly, Leu236Arg

4. Determination of the Quantities of Expression for BFP Mutants

Determination of the quantities of expression for the BFP mutants obtained is not particularly limited, but a comparison of the quantities of their

expression in *E. coli* by means of SDS-PAGE is preferable. Concretely, an overnight culture of *E. coli* into which each expression vector of pQE30 (empty vector), pQE30(201), and pQE30(202) had been introduced was diluted to 1/50 and it was grown in 3 ml of 2xYT carbenicillin medium at 37 °C for 3 h. IPTG was added to each sample to give its final concentration of 0.24 mg/ml, and the induction of a BFP protein was performed by further culturing the sample for 2.5 h.

An aliquot (100 μ l) was taken out from each sample and centrifuged, and precipitates were dissolved in a sample buffer. For each sample, 1.3 ml of *E. coli* was centrifuged at 10,000 rpm for 1 min and precipitates were suspended in 260 μ l of PBS(-). This suspension was frozen and thawed at -80 °C for 10 min, and was subjected to ultrasonic treatment (Elma Transonic ultrasonic washer 460/H). Subsequently, it was centrifuged at 15,000 rpm for 5 min to separate soluble proteins from insoluble fractions containing the inclusion body. These were subjected to SDS-PAGE in quantities that correspond to 50 μ l cultures of *E. coli* and were stained with Coomassie Brilliant Blue.

5. Comparison of Brightness of *E. coli* Cells Having a Variety of GFPs and BFPs Introduced

JM109 was transformed with each of pQE30 (empty vector), pQEGFP(101), pQEGFP(105), pQEBFP(201), pQEBFP(202), and pQEBFP(205), and it was streaked on a LB agar medium containing carbenicillin. After incubation at 37 °C for 24 h, the upper lid was removed and the plate was turned upside down and irradiated with UV (Funakoshi UV Transilluminator FTI-201 UV 365 nm to have photographs taken.

6. Transfection of GFP and BFP Mutant cDNAs into CHO Cells by the Calcium Phosphate Method and Fluorescence Measurements

A. Transfection

Coding regions were cut out from the pQE vectors containing the genes of GFP and BFP mutants that had been prepared by the site-directed mutation introduction method, and the corresponding portions of pHGFP(101)-C1 vectors were replaced by them; thus, pHGFP(103-105)-C1 and pHBFP(202-205)-C1 were prepared.

~~Unless otherwise so stated, CHO-K1 cells were grown in a F12+10% FBS medium in 5% CO₂ at 37 °C. The cells (1x10⁵) were inoculated into a 6-cm dish, and on the following day, their transfection was conducted in two dishes as a pair by the calcium phosphate method. (C. Chen and H. Okayama Mol. Cell. Biol. 7: 2745-2752 (1987).) After transfection, the one dish was incubated at 37 °C and the other at~~

INSAT
20

Q

~~30 °C for 24 h. The transfected CHO cells were~~
 a washed with 1xPBS(-) three times, and they were
~~dissolved in 1 ml of 10 mMTris-HCl (pH 7.4)~~
 containing 1% Triton X-100 and recovered in an
 5 Eppendorf tube. A supernatant (0.5 ml) from
 centrifugation at 3,000 rpm for 5 min was diluted 4-
 fold with the same buffer and fluorescence
 measurement was performed. Here, a pUC2SR α MCS
 vector (empty vector) was transfected and used as a
 10 blank. A Hitachi F-2000 type fluorophotometer was
 used in the fluorescence measurement. In the
 measurement of GFPs, fluorescence was scanned between
 460 nm and 600 nm at an excitation wavelength of 460
 nm to measure the maximal value in the vicinity of
 15 the fluorescence wavelength of 510 nm. In the
~~measurement of BFPs, fluorescence was scanned between~~
 a 360 nm and 500 nm at an excitation wavelength of 360
 nm to measure the maximal value in the vicinity of
~~the fluorescence wavelength of 445 nm.~~

20 7. Western Blotting

INSA8 ~~The CHO cells were transfected with pUC2SR α MCS~~
 1.08 (empty vector) (T. Tsukamoto et al. Nature Genet. 11:
 395-401 (1995)), pHGFP(101)-Cl, pHGFP(105)-Cl,
 pHBFP(201)-Cl, and pHBFP(205)-Cl, respectively and
 25 grown at 37 °C and at 30 °C. Employing a sample
 prior to dilution as used in the fluorescence

measurement previously described (8 //1), ~~SDS-PAGE~~
 was performed on a 12% gel. With the use of a
 Horizonblot (ATTO Inc.), transfer was conducted onto
 a nitrocellulose membrane (Millipore Inc., HAHY394FO)
 under the conditions of 2 mA and 90 min per cm².
 After the membrane was taken out and washed with
 1xPBS, it was immersed in 1% skim milk/PBS and shaken
 at room temperature for 30 min. After the membrane
 was washed with 1xPBS, it was immersed in 0.1% skim
 milk/PBS containing an anti-GFP antibody (Clonetech
 Inc.) that had been diluted 2,000-fold and shaken at
 4 °C overnight. The membrane was washed with 1xPBS
 for 5 min, and then with TPBS (0.05% Trion X-100/PBS)
 for 15 min three times. The membrane was immersed in
 0.1% skim milk/PBS containing an anti-rabbit IgG
 antibody labeled with HRP (Amersham Inc.) that had
 been diluted 1,000-fold, and shaken at 4 °C for 1 h.
 The membrane was washed with 1xPBS for 5 min, and
 then with TPBS (0.05% Trion X-100/PBS) for 15 min
 three times. The membrane was reacted with a
 chemiluminescence reagent (Amersham Inc. ECL) for 1
 min, and then, was exposed to an X-ray film for 2 min.

(II) Amino Acid Sequences of Novel GFP and BFP Mutants

1. Sequence Determination of BFP Mutants

Among the 10 mutants obtained, one mutant

(Mutant No. 10) proved that phenylalanine at amino acid number 64, which had been at the immediate N-terminal side of the chromophore, mutated into leucine.

5 With respect to this mutant clone, another mutation (L236R) had been introduced into its C-terminus (Table 1)

TABLE 1

10	mutant no. mutation
	1 L(CTT)1H(CAT)
	2 D(GAT)7Y(TAT)
	3 I(ATC)6T(ACC)
	4 the multicloning site: 14bp deletion
15	from BamHI
	5 the multicloning site: 24bp deletion
	from BamHI
	6 I(ATC)6N(AAC)
	7 L(CTT)4P(CCT), I(ATC)128G(GTC),
20	D(GAC)197A(GCC), S(AGC)202C(TGC)
	8 L(CTT)4R(CGT)
	9 M(ATG)1T(ACG), Y(TAC)39N(AAC), K(AAG)52E(GAG)
	10 K(AAG)41K(AAA)silent, F(TTC)64L(CTC),
	L(CTG)236R(CGG)
25	-----

With respect to this mutant, BFPcDNA was

subcloned into the same HindIII site as in pQEBFP(201) for a comparison purpose to prepare pQEBFP(202).

5 2. Comparison of the Quantities of Expression for BFP Mutants in *E. coli* by SDS-PAGE

10 IPTG was added to *E. coli* cultures harboring pQEBFP(201) and pQEBFP(202) and BFP proteins were allowed to express. When the *E. coli* cells were irradiated with UV, the *E. coli* harboring pQEBFP(202) apparently exhibited stronger fluorescence. When the proteins from these *E. coli* were analyzed by SDS-PAGE, the production of the 31 kDa protein was recognized to almost similar degrees in both *E. coli* having the respective plasmids (FIG. 1, Lanes 4 and 7).

15 When the solubility of these BFPs was also studied, BFP(201) with weaker fluorescence was nearly insoluble (FIG. 1, Lanes 5 and 6), whereas BFP(202) was mostly recovered in the soluble portion (FIG. 1, Lanes 8 and 9).

20 3. Comparison of Fluorescence of *E. coli* Cells Having a Variety of GFPs and BFPs Introduced

GFPs and BFPs into which the mutations of V163A and S175G had been further introduced in addition to F64L were prepared (see Table 4).

25 In order to compare the intensities of fluorescence in *E. coli*, streaking was performed

using *E. coli* cells having an empty pQE30 vector or pQE30 vectors into which cDNAs of GFP101, GFP105, BFP201, BFP202, and BFP205 had been subcloned. The *E. coli* having the empty vector introduced was not luminous. The *E. coli* having BFP201 prior to its improvement subcloned, even when irradiated with UV, was hardly luminous. In contrast, the one into which 202 had been subcloned was brightly luminous in blue. Further, it could be ascertained that 205 was even more brightly luminous than was 202.

As for GFPs, green fluorescence was observed by the naked eye, and a distinctive difference in brightness was noted between 101 and 105 (FIG. 2).

6. Transfection of GFP and BFP Mutant cDNAs into CHO Cells and Fluorescence Measurements

Since very luminous GFPs and BFPs were obtained in *E. coli*, the comparison was made also in mammalian cells (CHO). The results from the fluorescence measurements of cell extracts that were already prepared under culturing at 37 °C and at 30 °C are summarized (Table 5).

TABLE 5

	GFP or BFP	37 °C	30 °C
	101	30.8	214.6
5	103	532.1	765.4
	104	659.0	697.9
	105	2991.1	868.7
	201	14.3	166.7
	202	304.6	188.6
10	203	331.3	210.9
	204	330.9	265.9
	205	901.5	287.7

15 The values shown in the table are those obtained by subtracting the value of the empty vector used as a blank from the values of fluorescence obtained. The blank values were 8.9 in the measurement of GFPs at 37 °C, 7.14 in the measurement at 30 °C, 64.3 in the measurement of BFPs at 37 °C, and 50 in the measurement at 30 °C.

20 Table 6 shows relative values when the fluorescence intensity of GFP or BFP prior to its improvement after culturing at 37 °C is taken as 100, and it also makes comparisons in terms of ratio of fluorescence at 37 °C to that at 30 °C. From Table 6,

5 BFP(202) having the mutation as found by the
Mutagenic PCR exhibited the fluorescence 21 times
stronger at 37 °C. Further, BFP(202) had two
mutations (F64L and L236R); however, BFP(203) having
only F64L exhibited a similar intensity of
fluorescence to that of 202. This mutation is
believed to have caused stronger fluorescence.
Seventeen times stronger fluorescence was observed in
GFP(103) having F64L.

10 On the other hand, BFP(204) and GFP(104), both
of which had the mutations of V163A and S175G, were
brighter 23 times and 21 times, respectively.
GFP(105) and BFP(205) in which these mutations were
combined with F64L mutation were brighter 97 times
15 and 63 times. In addition, when the ratios of
fluorescence intensities at 37 °C to those at 30 °C
are taken for comparison, either of 101 and 201 prior
to its improvement was darker at 37 °C than at 30 °C.
Those having F64L alone or the combination of V163A
20 and S175G showed increases in the ratio of
fluorescence intensities at two temperatures, whereas
it was found that the fluorescence at 37 °C was more
than three times brighter with respect to GFP(105)
and BFP(205) in which the mutations were combined
25 (Table 6).

TABLE 6

	GFP or BFP	37 °C	37 °C/30 °C
	101	100	0.14
5	103	1728	0.70
	104	2140	0.94
	105	9711	3.44
	201	100	0.09
	202	2130	1.62
10	203	2317	1.57
	204	2314	1.24
	205	6304	3.13

15 6. Examination of the Quantities of Expression in
Animal Cells by Means of Western Blotting

20 The CHO cells were transfected with pUCD2SR α MCS
(empty vector), phGFP(101)-Cl, phGFP(105)-Cl,
phBFP(201)-Cl, and phBFP(205)-Cl, respectively and
cultured at 37 °C and 30 °C. Employing an anti-GFP
antibody for the cultured cells, the quantities of
GFP or BFP proteins expressed were examined. About
30kD bands that were not recognized in the
transfection of the empty vector (FIG. 3, Lanes 1 and
25 6) were detected.

In culturing at 30 °C, no distinctive difference

was noted between the content of GFP or BFP proteins expressed prior to the introduction of mutations and that after the introduction of mutations (Lanes 7-10). On the other hand, in culturing at 37 °C it was found that the mutants (Lanes 3 and 5) clearly expressed the GFP and BFP proteins in larger quantities (FIG. 3, Lanes 2-5).

The effects associated with the improved BFP and GFP mutants according to this invention are summarized below.

(1) The mutant type BFP(202) obtained by the Mutagenic PCR exhibits enhanced fluorescence as compared to BFP prior to the introduction of mutation in either *E. coli* cells or mammalian cells. In the clone of said mutant BFP, phenylalanine at amino acid number 64 has mutated into leucine (F64L), and further, leucine (amino acid number 236 at the C-terminus) has mutated into arginine (L236R).

With respect to the mutant type BFP(203) having only the mutation at amino acid number 64 as described above, a similar enhancement in fluorescence was also noted in mammalian cells. Therefore, it is F64L that is the responsible mutation for this mutant type BFP(202).

Such a mutation is presumed to involve a mechanism similar to the fluorescence enhancement

reported for GFP. (T. -T. Yang et al. Nucleic Acids Res. 24: 4592-4593 (1996).)

(2) The quantities of expression of proteins and the production of soluble proteins were investigated:

- 5 (i) Although the content of proteins is the same based on the comparison of the quantities of expression of Mutant BFPs in *E. coli* (through SDS-PAGE), the proteins from the mutant type BFP(201) are mostly insoluble whereas soluble proteins have
10 increased in the mutant type BFP(202); and (ii) a large difference in brightness was also seen in *E. coli*. These results indicate that the mutant type BFP(201) cannot correctly occupy a higher-order structure such as the formation of a chromophore
15 whereas the mutant type BFP(202) tends to occupy a more correct higher-order structure with ease: the mechanism for the above-mentioned fluorescence enhancement is believed to be due to this.

- 20 (3) On the other hand, the results of western blotting in the mammalian cells show that the quantity of proteins from GFP or BFP itself has increased. Namely, it is thought that the protein can occupy a stabilized higher-order structure in the mammalian cells; or alternatively, proteolysis
25 becomes slower than that prior to the improvement because the protein structure is stabilized.

(4) With the introduction of the F64L mutation having the characteristics as described above and other mutations, V163A and S175G, GFP and BFP proteins that have markedly improved characteristics in the expression at 37 °C in addition to those as described above are obtained.

Accordingly, the improved types of GFPs and BFPs into which such mutations have been introduced are provided with the characteristics that will allow them to be brightly luminous even at 37 °C, and they will enable observation in the mammalian cells where culturing is to be conducted at 37 °C. These improved types of GFPs and BFPs can be applied to cell biology as well as to many research areas. FIGS. 4(A) to (B) show the effects of this invention as described above.

From the invention thus described, it will be obvious that the invention may be varied in many ways. Such variations are not to be regarded as a departure from the spirit and scope of the invention, and all such modifications as would be obvious to one skilled in the art are intended for inclusion within the scope of the following claims.